PROTOCOL

05\2022

Materials:

- Chicken Forte



Method:

- 400 ml (explained in use instruction) of Chicken Forte for every 1,000 litres of drinking water for the chickens.
- The product should be applied from day 1.
- A water consumption per chicken of 9 litres in 6 weeks (42 days) is estimated.
- Thus, the consumption of Chicken Forte every 1,000 chickens is 3.6 liters in 42 days of the life of the chickens.

Use instructions:

- Our product is sterile, before you begin with our product, be sure that your water pipes are sterile and clean.
- Please make sure the water tank before the pump is clean and sterile.
- Be sure that you put the right concentration in the pump, please feel free to contact us to any technical advice of any equipment you have.
- For high consumption of the product, you must pick one of two methods:

  1) use 400 ml of chicken forte, and every four days you must clean the water drinking pipeline by inserting a water flow for 2 to 3 minutes and draining out the residual matter from the end of the pipes.
  Or if you have an automatic pipeline, you can clean the pipeline automatically with water flow (disinfectant is preferred).

Natura Feed ltd, Industrial zone, Eilaboun, Israel, Postal code: 1697200, P.O.B 442, Tel: +97246780302
Fax: +97246780303, office@naturafeedpro.com, www.naturafeedpro.com



2) In case of any infection or disease – Raise the Dosage to 600 ml for three continuous days.

- Please from now and then, check the end of the pipes for any residual matter. If found, please insert a water flow while drawing out water from the end of the drinking line. Use disinfectant if needed.

Total Usage of Chicken Forte:

- 25,000 broilers * 9 Liter of water * 400 ml /1000 L = ~ 90 Liters of Chicken Forte for 42 days.

Control:

- Nearby Coop with 25,000 Live mixed chickens (Not treated with Chicken Forte): same receiving day, same ancestor, same chicken house conditions.

The Parameters\data collecting:

- Parameter that must be received in the first trial with chicken forte:

	1 day old	Two weeks	21 days	28 days	35 days	42 days	Slaughter house
Broiler/m^2	√	√	√	√	√	√	
Weight 50 males	Average	√	√	√	√	√	Truck √
Weight 50 females	√	√	√	√	√	√	
% mortality		√	√	√	√	√	√
FCR		√	√	√	√	√	√
European index						At the end √	
Meat/m^2							√
% yield of dressing percentage							√